

- JONES, G. N. 1950. Flora of Illinois, ed. 2. Am. Midl. Nat. Monogr. 5: 1-368. University of Notre Dame Press, Notre Dame, Indiana.
- \_\_\_\_\_. 1963. Flora of Illinois, ed. 3. Am. Midl. Nat. Monogr. 7: 1-401. University of Notre Dame Press, Notre Dame, Indiana.
- PATTERSON, H. N. 1867. Catalogue of phaenogamous and vascular cryptogamous plants of Illinois. Oquawka, Illinois. 54 pp.
- STEYERMARK, J. A. 1963. Flora of Missouri. The Iowa State University Press, Ames, Iowa. 1725 pp.
- SVENSON, H. K. 1957. *Fuirena*. In: North American Flora 18(9): 505-507.

SOUTHERN ILLINOIS UNIVERSITY  
CARBONDALE, ILLINOIS 62901

---

## TWO NEW COLOR-FORMS FROM SOUTHERN MAINE

While collecting in the Ocean Park section of Old Orchard Beach, York County, Maine, in recent summers, I have noticed certain plants of *Spiraea latifolia*, Meadowsweet, with deep-pink flowers. Some of these plants have pubescent panicles; others are glabrous. I have found this form growing both in open wet spots and in shaded woodland stations.

*Spiraea latifolia* (Ait.) Borkh. forma *rosea* T. W. Wells, forma nova. TYPE: Maine: YORK CO.: Old Orchard Beach, 11 October, 1969, T. W. WELLS (NEBC). Differt a forma *latifolia* floribus valde roseis.

While collecting in salt-marshes along Goose Fare Creek, which is the southern boundary of Ocean Park, I found plants of *Polygonum ramosissimum* which differ from the typical in that the sepals have bright-white margins.

*Polygonum ramosissimum* Michx. forma *alba* T. W. Wells, forma nova. TYPE: Maine: YORK CO.: Old Orchard Beach, 5 September, 1970 T. W. WELLS (NEBC). Differt a forma *ramosissimum* sepalis albomarginatis.

THEODORE W. WELLS  
MILTON ACADEMY  
MILTON, MASSACHUSETTS 02186